

The Linear $U3 \times U3$ Sigma Model, the $\sigma(\approx 500)$ and the Spontaneous Breaking of Symmetries¹

Nils A. Törnqvist

Physics Department, POB 9, FIN-00014, University of Helsinki, Finland

Abstract. My recent fit of the light scalar mesons is discussed in the light of the $U3 \times U3$ linear sigma model. The argument for why there exists a light and broad $\sigma(\approx 500)$ meson is explained in more detail, and the mechanism of spontaneous symmetry breaking is discussed.

The first thing I want to point out is that my recent model fit to the light scalar $q\bar{q}$ nonet [1], in fact, can be interpreted as a model unitarization based upon the $U(3) \times U(3)$ chiral symmetric renormalizable Lagrangian

$$\mathcal{L} = \frac{1}{2}\text{Tr}[\partial_\mu \Sigma \partial_\mu \Sigma^\dagger] + \frac{1}{2}m^2\text{Tr}[\Sigma \Sigma^\dagger] - \lambda\text{Tr}[\Sigma \Sigma^\dagger \Sigma \Sigma^\dagger] - \lambda'(\text{Tr}[\Sigma \Sigma^\dagger])^2 + \mathcal{L}^{SB} . \quad (1)$$

Here $\Sigma = \sum_{i=0}^8 (s_i + ip_i)\lambda_i/\sqrt{2}$ are 3×3 matrices, s_i and p_i stand for the 0^{++} and 0^{-+} nonets and λ_i are the Gell-Mann matrices, normalized as $\text{Tr}[\lambda_i \lambda_j] = 2\delta_{ij}$, and where for the singlet $\lambda_0 = \sqrt{\frac{2}{3}}\mathbf{1}$ is included. Note that each meson from the start has a definite $SU3$ symmetry content, which in the quark model means that it has a definite $q\bar{q}$ content.

Apart from the term $\mathcal{L}^{SB}$, Eq.(1) is clearly invariant under $\Sigma \rightarrow U_L \Sigma U_R^\dagger$. In Eq. (1) I have contrary to the usual convention defined the sign of m^2 such that the naive physical squared mass would be $-m^2$, and the instability thus occurs when $m^2 > 0$. Generally one includes in addition into $\mathcal{L}$ terms which break the symmetries:

$$\mathcal{L}^{SB} = c \cdot [\det \Sigma + \det \Sigma^\dagger] + \epsilon_0 s_0 + \epsilon_8 s_8 . \quad (2)$$

The c term breaks the $U(1)$ symmetry explicitly and gives the η_1 a mass, while ϵ_0 gives the pseudoscalar octet mass and ϵ_8 breaks explicitly $SU3_f$ down to isospin.

¹⁾ Talk given at Hadron97 or International Conference on Hadron Spectroscopy, Brookhaven, USA Aug 24-30 1997. And at the EuroDaphne meeting in Barcelona Nov. 6-9 1997.

As well known with $-m^2 < 0$ and $\lambda > 0, \lambda' > 0$ the potential in Eq.(1) (being of the form of a "Mexican hat") gives rise to an instability with vacuum condensate $\langle s_0 \rangle = v = \sqrt{(2/3)}f_\pi$. Let $\lambda' = 0$ for a moment, then shifting as usual the scalar field, $\Sigma \rightarrow \Sigma + v$, one finds $v = m^2/(4\lambda)$. Furthermore, the squared mass ($-m^2$) is for the 0^{++} nonet replaced by $-m^2 + 12\lambda v^2 = 2m^2$, while the 0^{-+} nonet becomes massless, $-m^2 + 4\lambda v = 0$. If we had included the λ' term instead of the λ term, then only the scalar singlet would acquire mass while the remaining 17 states would be massless.

In addition, very importantly, after shifting the scalar singlet field, the λ term (keeping $\lambda' = 0$) generates trilinear spp and sss couplings of the form $g[\text{Tr}[\lambda_i \lambda_j \lambda_k + h.c.]]/2$ and with an overall normalization $g = 4\lambda f_\pi$. These couplings obey the simply connected, Okubo-Zweig-Iizuka (OZI) allowed quark line rules with flavour symmetry exact. One has $\text{SU}3_f$ predictions relating different couplings constants. Denoting by σ the $u\bar{u} + d\bar{d}$ scalar, and by σ_s the $s\bar{s}$ scalar one has e.g. $g^2 = g_{\sigma\pi^0\pi^0}^2 = \frac{1}{2}g_{\sigma\pi^+\pi^-}^2 = g_{\sigma K\bar{K}}^2 = \frac{2}{3}g_{K_0^*K\pi}^2 = g_{a_0K\bar{K}}^2 = \frac{1}{2}g_{\sigma_s K\bar{K}}^2$, $g_{\sigma_s\pi\pi} = 0$ etc. Here we summed over charge states except for the $\sigma\pi\pi$ couplings since conventionally the $\sigma\pi\pi$ coupling is $g_{\sigma\pi^0\pi^0}$. If one includes also the λ' term of Eq.(1) then only the couplings involving the σ and σ_s states would be altered, which would violate the OZI rule at the tree level.

Such flavour symmetric OZI couplings, generated by the λ term of Eq.(1), together with a near-degenerate bare scalar nonet mass were, in fact, the starting point of my recent analysis of the scalar $q\bar{q}$ nonet [1]. In particular it was crucial that after determining the overall coupling g from a fit to data on $K_0^* \rightarrow K\pi$ and $a_0 \rightarrow \pi\eta$ one predicted correctly the $\pi\pi$ phase shifts. This shows that the above relations relating the bare σ and σ_s couplings to the same overall g as those of $K_0^*K\pi$ and $a_0\pi\eta$ must be approximately satisfied experimentally. I.e., one cannot tolerate very big λ' coupling in Eq.(1) since then these relations would be destroyed. Another argument for that λ' must be small is that then the bare scalar singlet state would have a very different mass from the other nonet members, not needed in Ref. [1].

After the unitarization was performed in [1] the scalars acquire finite widths and are strongly shifted in mass by the different couplings to the $0^{-+}0^{-+}$ thresholds. The 0^{-+} masses in the thresholds were given their experimental values, and consequently the main source of flavour symmetry breaking in the output physical mass spectrum was generated by the vastly different positions of these thresholds. E.g. the large experimental splitting between the $a_0(980)$ and $K_0^*(1430)$ masses came from the large breaking in the sum of loops for the K_0^* ($K_0^* \rightarrow K\pi, K\eta, K\eta' \rightarrow K_0^*$) compared to those for a_0 ($a_0 \rightarrow \pi\eta, K\bar{K}, \pi\eta' \rightarrow a_0$), although in the strict $\text{SU}3_f$ limit these loops give the same shift to the two resonances.

There were only 6 parameters in [1] out of which two parametrized the bare scalar spectrum (1.42 GeV for the $u\bar{u}, d\bar{d}, u\bar{d}, d\bar{u}$, and an extra 0.1 GeV when an s quark replaces a u or d quark). The overall coupling was parametrized by $\gamma = 1.14$ and $k_0 = 0.56$ GeV/c was the cut off parameter. Now, the γ parameter can be

related to λ of Eq.(1) through $\lambda = 2\pi\gamma^2 = 8.17$, by comparing the $\sigma\pi\pi$ coupling of the two schemes. [One has $g_{\sigma\pi^0\pi^0}^2/(4\pi) = \gamma^2 m_\sigma^2 = \lambda m_\sigma^2/(2\pi)$. The latter equation follows from $g_{\sigma\pi^0\pi^0} = g = 4v\lambda$ and $v^2 = -m^2/(4\lambda) = m_\sigma^2/(8\lambda)$ given above.]

Using the conventional value for $f_\pi = \sqrt{(2/3)}v = 93$ MeV and $\lambda = 8.17$ one predicts from (1) at the tree-level that the average mass of the $u\bar{u}$, $d\bar{d}$, $d\bar{u}$ states is $m = v(8\lambda)^{1/2} = 920$ MeV. Now although strictly speaking there is no exact one-one correspondence between the model of Eq. (1) at the tree level, before renormalization, and the unitarised model of [1], it is remarkable that this prediction is close to the average of the σ and $a_0(980)$ masses found in [1]. I.e. if we had used $f_\pi = 93$ MeV to determine the energy scale, one could have eliminated one of the 6 parameters.




FIGURE 1. The full line curve shows the $I=0$, phase shifts for $\pi\pi \rightarrow \pi\pi$, as predicted by the model [1]. If one lets the $s\bar{s}$ meson mass become very large (10 GeV) in the model then the rapid rise at 1 GeV due to the $f_0(980)$ vanishes. This shows that $f_0(980)$ is the unitarized $s\bar{s}$ state. The remaining phase shift (dot-dashed thick line) is then entirely due to the $u\bar{u} + d\bar{d}$ channel and one sees that what remains can be understood as a very broad resonance. This is the " σ ", whose 90° mass value is at 880 MeV, while the pole is at $470-i250$ MeV. (The thin dot-dashed curve shows the phase shift if also the $K\bar{K}$ and $\eta\eta$ thresholds are given large values.)

Thus an important point which I did not realize when writing [1] is that that model can be interpreted as an effective field theory given by $\mathcal{L}$ of Eq.(1) with $\lambda = 8.14$ and $\lambda' \approx 0$. The absence of the λ' term at the tree level means that the OZI rule holds exactly before unitarization. The three small additional terms in

$\mathcal{L}^{SB}$ in Eq.(2) give the pseudoscalars their physical masses, and account for the extra 0.1 GeV for the bare strange states.

Of course the unitarization procedure of [1] should be improved upon, including u - and t -channel singularities [2], more loops etc., but I believe that the dominant effects were already included phenomenologically for the scalar states.

An important point observed in the second paper of Ref. [1] was that the model requires the existence of the light and broad σ resonance. This is explained in more detail in Fig.1 above. Both before and at this conference there has appeared several new papers [3], which through different analyses and models support this same conclusion, i.e. that the light and broad sigma, which has been controversial for so long, really exists, and is here to stay [4].

Finally I have too little time left to explain sufficiently another recent argument of mine [5], which certainly many people would consider speculative. It involves the possibility that the spontaneous breaking of symmetries does not stop after the standard step discussed above. After quantum loops further symmetry breaking can occur, but here I refer interested to [5] and papers in progress.

REFERENCES

1. N. A. Törnqvist, Z. Physik **C68** 647 (1995); N. A. Törnqvist and M. Roos, Phys. Rev. Lett. **76** (1996) 1575.
2. N. Isgur and J. Speth, Phys. Rev. Lett. (Comment) **77** (1996) 2332; N.A, Törnqvist and M. Roos, *ibid* **77** (1996) 2333.
3. J. A. Oller and E. Oset, these proceedings (hep-ph/971554) and Nucl. Phys. **A620** (1997) 438;
 See the five different talks in these proceedings given by M.Y Ischida, S. Ischida, T. Ischida, and K. Takamatsu, and S. Ischida et al. Prog. Theor. Phys. **95** (1995) 745;
 See talk by R. Kaminski and Phys. Rev. **D50** (1994) 3145;
 L.R. Baboukhadia et al., J. Phys. **G23** (1997) 1065;
 A., Bramon et al. "Double counting ambiguities in the linear sigma model" (J. Phys. C to appear), hep-ph/9709274;
 M. Harada et al. Phys. Rev. Lett. (Comment) **78** (1997) 1603; N.A, Törnqvist and M. Roos, *ibid* **78** (1997) 1604;
 M.D. Scadron, "Scalar σ meson via chiral and crossing dynamics" hep-ph/9710317;
 See also M. Boglione and M. Pennington, Phys. Rev. Lett. **79** (1997) 1998.
4. R. M. Barnett et al., (The Particle Data Group), Phys. Rev. **D54** (1996) 1.
5. N.A. Törnqvist, Phys. Lett. **B406** (1997) 70.